

A Summary and Perspective of Recent IRIS Assessments and Impact on Planning Environmental Investigations

Anita K. Meyer DABT

Environmental & Munitions Center of Expertise

US Army Engineering and Support Center, Huntsville

EDQM Workshop

Arlington, VA

30 March 2011



BUILDING STRONG®

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 MAR 2011		2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE A Summary and Perspective of Recent IRIS Assessments and Impact on Planning Environmental Investigations				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Engineering and Support Center,Environmental & Munitions Center of Expertise,PO Box 1600,Huntsville,AL,35807				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Presented at the 2011 DoD Environmental Monitoring & Data Quality Workshop (EMDQ 2011), 28 Mar ? 1 Apr, Arlington, VA.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 15	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Role of Risk Assessment on Environmental Projects

- Used to determine whether site requires further study or remediation
- Used with regulatory values to determine cleanup levels
- Risk-based screening levels (aka PRGs, RBCs, RSLs)
 - ▶ Screen sites early in project lifecycle
 - ▶ Determine project quantitation limits



Risk-Based Values Function of Toxicity and Exposure

$$\text{Risk} = \frac{\text{Intake}}{\text{Toxicity}}$$

↑ Toxicity = ↓ Criteria

- Tiered approach used to identify toxicity values for site risk assessments
- Integrated Risk Information System (IRIS)
- IRIS values also inform regulatory decisions (MCLs etc.)



Emerging Contaminants (ECs)


- Are chemicals or materials of interest that are characterized by:
 - ▶ a perceived or real threat to human health or environment, and
 - ▶ there is no currently published health standard or there is an existing health standard, but the standard is evolving or being re-evaluated.

Source: “Initiation of Emerging Contaminants Characterization and Response Actions for Protection of Human Health” Issue Paper (ECOS & DoD Sustainability Workgroup, 2008)



DoD Participation in IRIS Interagency Reviews

Assessment Development Process for New IRIS



Tetrachloroethylene

Status: External Peer Review

- 1998 initiated
- June 2008 external review version released
- Nat'l Academy Review Feb 2006
- Current external (SAB) and public review

Risk-Based Screening Levels*			
	Res. Soil (mg/kg)	Res. Water Use (µg/L)	Indoor Air (µg/m ³)
Current	0.55	0.11	0.41
New (draft)	0.293	0.179	0.122
Sources of current toxicity values include EPA IRIS, ATSDR and CalEPA. Lowest RSL target risk = 10^{-6} . Draft values not suitable for project use.			



Trichloroethylene

Status: External Peer Review

- IRIS values withdrawn late 1980s
 - 2001 draft released
 - 2006 NAS review
 - 2009 re-released
 - Lowest RSLs based on 10^{-6} cancer risk
- Do not utilize levels derived from draft toxicity values.


Risk-Based Screening Levels*

	Res. Soil (mg/kg)	Res. Water Use (µg/L)	Indoor Air (µg/m ³)
Current	2.8	2	1.2
New (draft)	0.48	0.23	0.24
CalEPA source of current values. Lowest RSL target risk = 10^{-6} . Draft values not suitable for project use.			



Risk-Based TCE Residential Indoor Air Concentrations

1E-04 to 1E-06 cancer risk and Noncancer



Dioxin

Status: External Peer Review

- Assessment initiated in 1990
- Released for public and peer review 2010
- Using cancer toxicity values in this draft residential screening levels could be as low as 0.45 ppt dioxin toxicity equivalent (TEQ, for dioxin-like compounds)
- Current EPA policy recommends 1000 ppt
 - ▶ Interim PRG may be released



Final IRIS Reference Dose for Cis-1,2-Dichloroethylene (September 2010)

- Oral noncancer reference dose (RfD) = 0.002 mg/kg-day; increased kidney weight
 - ▶ PPRTV used until this time
 - ▶ MCL = 70 µg/L
- No cancer toxicity values published

Risk-Based Screening Levels*		
	Residential Soil (mg/kg)	Residential Water Use (µg/L)
Old	780	370
New	156	73
Fold Change	5x	



*Using EPA Regional Screening Level Calculator

Final IRIS Reference Dose for Trans-1,2- Dichloroethylene (September 2010)

- Oral noncancer reference dose (RfD) = 0.02 mg/kg-day;
- Based on decreased antibody production by the spleen
- No change in RfD value, but change of critical effect
- MCL = 100 µg/L
- No cancer toxicity values published



Final IRIS Values for 1,4- Dioxane (August 2010)

- Oral noncancer reference dose (RfD) = 0.03 mg/kg-day; Liver & kidney effects
 - ▶ Inhalation RfC will be developed in separate document
- Oral cancer slope factor (SF) = 0.1 mg/kg-day⁻¹; Rodent liver tumors

Risk-Based Screening Levels*		
	Residential Soil (mg/kg)	Residential Water Use (µg/L)
Old	44.1	6.1
New	4.85	0.67
Fold Change	9x	



*Using EPA Regional Screening Level Calculator

BUILDING STRONG[®]

Final IRIS Values for Hydrogen Cyanide and Cyanide Salts (September 2010)

- New Oral noncancer RfD = 0.0006 mg/kg-day; male reproduction effects
- New inhalation RfC = 0.0008 mg/m³; effects on nervous system and thyroid
- RfD is 33 times more conservative than previous IRIS value
- RfC is 3.75 times more conservative than previous IRIS value




Basis for Changes

- New study data
 - ▶ Sensitive measurements
- Science policy
 - ▶ Mutagenic mode action adjustment
 - ▶ Uncertainty factor application



High-Throughput Screening Assays

batch testing of chemicals for pharmacological/toxicological endpoints using automated liquid handling, detectors, and data acquisition



The future!
EPA Next Generation Risk Assessment
New technology will lead to greater capacity and speed. Interpretation of results is a challenge; studies underway to link known effects with screening assay data.

Systems Exposure Science : Extending Network Analysis

Consider coupled networks spanning multiple levels of biological organization

